



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : G01N 33/53, 33/543, 33/577	A1	(11) International Publication Number: WO 85/ 02258 (43) International Publication Date: 23 May 1985 (23.05.85)
(21) International Application Number: PCT/US84/01888 (22) International Filing Date: 16 November 1984 (16.11.84) (31) Priority Application Number: 553,219 (32) Priority Date: 18 November 1983 (18.11.83) (33) Priority Country: US (71) Applicant: BECKMAN INSTRUMENTS, INC. [US/ US]; 2500 Harbor Boulevard, Fullerton, CA 92634 (US). (72) Inventors: CRAGLE, Linda, Katherine ; 4949 Cadden Way, San Diego, CA 92117 (US). HARRIS, Paul, C. ; 4740 Calle Bonita, Yorba Linda, CA 92686 (US). LEE, Shih-Yun ; 1311 McClelland Street, San Mar- cos, CA 92069 (US). TUNG, Ker, Kong ; 7739 Anillo Way, Carlsbad, CA 92008 (US). VODIAN, Morton, Alan ; 1770 Sally Place, Escondido, CA 92026 (US).	(74) Agents: MAY, William, H. et al.; Beckman Instru- ments, Inc., Legal Department, 2500 Harbor Boule- vard, Fullerton, CA 92634 (US). (81) Designated States: BE (European patent), DE (Euro- pean patent), FR (European patent), GB (European patent), JP, NL (European patent), SE (European pa- tent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: NOVEL IMMUNOASSAYS AND KITS FOR USE THEREIN

(57) Abstract

A improved immunoassay for assaying an antigenic substance (Ag) in a fluid. The immunoassay is the type which comprises contacting the fluid with at least one first entity selected from a group consisting of an antibody (Ab) to the Ag, a soluble, labeled antibody (L-Ab_a) to the Ag, and an antibody (Ab_b) to the Ag bound to a solid support (SC). The immunoassay is characterized in that the fluid is contacted with at least one additional entity selected from a group consisting of at least one different type of soluble, labeled antibody (L-Ab_c) to the Ag, at least one different type of antibody (Ab_d) bound to a solid carrier (SC₁), and at least one different type of antibody (Ab_e) to the Ag. The SC₁ is selected from a group consisting of SC, at least one different solid carrier (SC₂), and mixtures thereof (SC and SC₂). Each type of L-Ab_c, Ab_d-SC₁, and Ab_e has a lower average affinity constant (K) for Ag than each respective K of L-Ab_a, Ab_b-SC and Ab_e; and the additional entity is present in an amount sufficient to avoid a hook effect. Also, an improved reagent of the type comprising at least one first entity selected from a group consisting of L-Ab_a, Ab_b-SC, and Ab. The reagent is characterized in that it further comprises at least one additional entity selected from a group consisting of at least one different type of L-Ab_c and at least one different type Ab_d-SC₁, and at least one different type of Ab_e. Ab, L-Ab_a, Ab_b-SC, L-Ab_c, Ab_d-SC₁, and Ab_e are as defined above.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GA	Gabon	MR	Mauritania
AU	Australia	GB	United Kingdom	MW	Malawi
BB	Barbados	HU	Hungary	NL	Netherlands
BE	Belgium	IT	Italy	NO	Norway
BG	Bulgaria	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union -
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali		
FR	France				

NOVEL IMMUNOASSAYS
AND KITS FOR USE THEREIN

Background of the Invention

1. Field of the Invention

This invention relates to immunoassays and to kits for use therein.

2. Description of the Prior Art

Numberous types of immunoassays presently exist. To illustrate, one- and two-step sandwich immunoassays for antigens which combine to at least two antibodies are known (1-4). For example, U.S. Patent 4,244,940 (2) discloses a sandwich immunoassay method in which a sample containing an antigen (Ag) to be determined, a labeled antibody (L-Ab_a) to the Ag, an antibody (Ab_b) for the Ag bound to a solid-phase support (SC) are brought together in a single incubation mode or step in an aqueous medium to form a substantially stable suspension and produce a two-phase system. The solid phase portion of this two-phase system contains the Ab_b-SC, a portion of which has become bound to the Ag which in turn has become bound to a portion of the L-Ab_a (represented as L-Ab_a-Ag-Ab_b-SC). The liquid phase portion of the two-phase systems contains the unbound portion of the L-Ab_a. The solid and liquid phases are separated and either phase analyzed for the L-Ab_a, the concentration of which is a function of the concentration of Ag in the sample. U.S. Patent 4,244,940 (2) teaches that this two-site immunoassay employing a single incubation mode or step provides significant advantages over assay procedures involving more than one incubation by simplifying, shortening, and rendering more convenient the performance of the assay. Furthermore, U.S. Patent



4,244,940 (2) teaches that this improvement in assay procedure is accomplished while maintaining acceptable assay characteristics such as precision, specificity, and stability, in addition to being less subject to errors in timing, additions, and other manipulations.

One- and two-step sandwich immunoassays, can be used for the determination of the presence or concentration of any Ag which can simultaneously become bound by two antibodies. This group of Ags includes, but is not limited to, placental, pituitary, calcium regulating, and adrenal medullary polypeptide hormones, protein and protein fragments; immunoglobulins (antibodies) of various classes; viral, viral subunits, bacterial, and protozoal organisms or particles; toxins; drugs, enzymes, and tumor-associated antigens.

In the one- and two-step sandwich immunoassays, the antibody employed is any substance which binds the Ag with acceptable specificity and affinity.

In one- and two-step sandwich immunoassay the L-Ab_a may be labeled with any of a number of known tracers. These tracers include, but are not limited to, radioactive tags, fluorescent labels, and enzyme labels.

An important component of the one- and two-step sandwich immunoassays is the SC for the Ab_b. The SC must be able to (a) be bound to the Ab_b, (b) be handled conveniently during manipulations such as pipetting and centrifuging, and (c) exhibit low nonspecific adsorption properties or be treated so that it exhibits such adsorption properties.

A pitfall in both the conventional and one-step sandwich immunoassay has been reported (3,5-7). More

particularly, these authors have found that a potential hazard of misinterpreting the results is involved in both the conventional and one-step sandwich immunoassays because of a false negative result or "hook effect" at high concentrations of Ag.

Nomura et al. (6), who discuss this pitfall in regard to a one-step sandwich immunoassay with monoclonal antibodies for the determination of human alpha-fetoprotein (AFP), state that "theoretically but not practically, the inhibition in antigen excess region may be avoided by employing a large amount of immobilized and labeled anti-AFP." One reason that this suggestion is not practical is that the additional large amount of labeled anti-AFP suggested by Nomura et al. would result in an assay having high non-specific adsorption since non-specific adsorption is proportional to the concentration of L-Ab₂ employed. Another reason that this suggestion is not practical is that this additional large amount of labeled anti-AFP would reduce the dynamic range of the assay when an enzyme or other label requiring the use of a spectrophotometer is employed.

Miles et al. (7), who discuss this pitfall in regard to a two-step sandwich immunoassay, note that repeated washings after the first incubation does prevent the high dose hook effect. However, this technique for preventing the high dose hook effect is tedious and time consuming.

To date, there has been no convenient technique proposed for either preventing or avoiding this false negative or hook effect phenomenon in either one- or two-step sandwich immunoassays.

Another immunoassay known to those skilled in the art is the direct nephelometric immunoassay. The direct nephelometric immunoassay comprises contacting a fluid containing an antigenic substance (Ag) with an antibody (Ab) to the Ag in order to form a complex Ab-Ag. A measurement of the amount of formed Ab-Ag is directly proportional to the amount of Ag in the assayed fluid.

Like the one- and two-step sandwich immunoassays, there is a potential hazard of misinterpreting the results because of the false negative result or "hook effect" at high concentrations of Ag.

Although in the case of the direct nephelometric immunoassay there are some electromechanical methods for safeguarding against such misinterpretation, at present there is no known chemical means for either preventing or avoiding this false negative or hook effect phenomenon in this case either.

Accordingly, it would be very desirable to have immunoassay methodologies and kits wherein this problem has been either prevented or avoided.

Summary of the Invention

In accordance with the present invention, there are provided improved immunoassay methodologies and kits for use therein wherein all the advantages possessed by the prior immunoassays are maintained with the concurrent avoidance of the hook effect.

In general, the present invention encompasses an improved immunoassay for an antigenic substance (Ag) in a fluid. The immunoassay of the present invention is of the type which comprises contacting the fluid with at

least one first entity selected from a group consisting of an antibody (Ab) to the Ag, a soluble, labeled antibody (L-Ab_a) to the Ag, and an antibody (Ab_b) to the Ag bound to a solid carrier (SC). The present immunoassay is characterized in that the fluid is contacted with at least one additional entity selected from a group consisting of at least one different type of soluble, labeled antibody (L-Ab_c) to the Ag, at least one different type of antibody (Ab_d) to the Ag bound to a solid carrier (SC₁), and at least one different type of antibody (Ab_e) to the Ag.

SC₁ is selected from a group consisting of SC, one or more different solid carriers (SC₂), and mixtures thereof (SC and SC₂).

Each of the additional entities has an average affinity constant (K) for the Ag lower than the K of its corresponding first entity for the Ag. In addition, the additional entity is present in an amount sufficient to avoid the hook effect.

More particularly, the present invention encompasses an improved one-step sandwich immunoassay. This one-step sandwich immunoassay of the present invention is of the type which comprises:

- (a) contacting the fluid with (i) the L-Ab_a and (ii) the Ab_b bound to the SC to form an insoluble complex (L-Ab_a-Ag-Ab_b-SC);
- (b) separating the L-Ab_a-Ag-Ab_b-SC from the fluid and unreacted L-Ab_a; and

- (c) measuring either the amount of $L\text{-}Ab_a$ associated with the $L\text{-}Ab_a\text{-}Ag\text{-}Ab_b\text{-}SC$ or the amount of unreacted $L\text{-}Ab_a$.

The improved one-step sandwich immunoassay of this invention is characterized in that the fluid is contacted with at least one additional entity selected from the group consisting of one or more different type of $L\text{-}Ab_c$ and one or more different type of Ab_d bound to the SC_1 ;

wherein:

- (i) each different type of $L\text{-}Ab_c$ and $Ab_d\text{-}SC_1$ has a lower K for Ag than each respective K of $L\text{-}Ab_a$ and $Ab_b\text{-}SC$ for Ag ; and
- (ii) the additional entity is present in an amount sufficient to avoid the hook effect.

More particularly, the present invention also encompasses an improved two-step sandwich immunoassay. This immunoassay is of the type which comprises:

- (a) contacting the fluid containing the Ag with the $Ab_b\text{-}SC$ to form an insoluble complex ($Ag\text{-}Ab_b\text{-}SC$);
- (b) contacting the $Ag\text{-}Ab\text{-}SC$ with the $L\text{-}Ab_a$ to form an extended insoluble complex ($L\text{-}Ab_a\text{-}Ag\text{-}Ab_b\text{-}SC$);
- (c) separating the $L\text{-}Ab_a\text{-}Ag\text{-}Ab_b\text{-}SC$ from unreacted $L\text{-}Ab_a$; and
- (d) measuring either the amount of $L\text{-}Ab_a$ associated with the $L\text{-}Ab_a\text{-}Ag\text{-}Ab_b\text{-}SC$ or the amount of unreacted $L\text{-}Ab_a$.

The improved two-step sandwich immunoassay of this invention is characterized in that in step (b) the $\text{Ag-Ab}_b\text{-SC}$ is contacted with one or more different type of L-Ab_c , wherein each different type of L-Ab_c has a lower K for Ag than the K of L-Ab_a and L-Ab_c is present in an amount sufficient to avoid the hook effect.

More particularly, the present invention also encompasses an improved direct nephelometric immunoassay. This immunoassay is of the type which comprises:

- (a) contacting the fluid with Ab in order to form a complex Ab-Ag ; and
- (b) measuring the amount of formed Ab-Ag .

The improved direct nephelometric immunoassay of this invention is characterized in that the fluid is contacted with at least one additional type of antibody Ab_e . Each type of Ab_e has a K for the Ag lower than the K of the Ab for the Ag and is present in an amount sufficient to avoid the hook effect.

Also within the general scope of this invention is an improved reagent. This improved reagent is of the type comprising at least one first entity selected from a group consisting of Ab , L-Ab_a , and $\text{Ab}_b\text{-SC}$. The reagent of the present invention is characterized in that it further comprises at least one additional entity selected from a group consisting of at least one different type of L-Ab_c , at least one different type of $\text{Ab}_d\text{-SC}_1$, and at least one different type of Ab_e . The additional entity is present in an amount sufficient to avoid a hook effect when the reagent is employed in an immunoassay.

In the case of one- and two- step sandwich immunoassays, the reagent is of the type comprising $L\text{-Ab}_a$ and $\text{Ab}_b\text{-SC}$. The improved reagent of the present invention is characterized in that it further comprises at least one additional entity selected from a group consisting of one or more different types of $L\text{-Ab}_c$ and one or more different types of Ab_d bound to SC_1 . The additional entity is present in an amount sufficient to avoid the hook effect when the reagent is employed in the immunoassay.

In the case of direct nephelometric immunoassays, the reagent is of the type comprising Ab . The improved reagent of the present invention further comprises at least one different type of Ab_e in an amount sufficient to avoid the hook effect when employed in a direct nephelometric assay.

Description of the Preferred Embodiments

The average affinity constant (K) for each different type of $L\text{-Ab}_c$, $\text{Ab}_d\text{-SC}_1$, and Ab_e for Ag must be lower than each respective K of $L\text{-Ab}_a$, $\text{Ab}_b\text{-SC}$ and Ab for Ag . This restriction assures that the former entities compete at most only insignificantly with the latter entities in the dynamic range. In order to achieve this result, it is preferred that each K of each different type of $L\text{-Ab}_c$, $\text{Ab}_d\text{-SC}_1$, and Ab_e for Ag be at least 5, more preferably at least 10 times lower than each respective K of $L\text{-Ab}_a$, $\text{Ab}_b\text{-SC}$, and Ab for Ag .

In addition, the further one wishes to extend the avoidance of the hook effect, the larger the difference must be between (a) the K of $L\text{-Ab}_a$ and the K of $L\text{-Ab}_c$, (b) the K of $\text{Ab}_b\text{-SC}$ and the K of $\text{Ab}_d\text{-SC}_1$, and (c) the K of Ab and the K of Ab_e .

When more than one type of L-Ab_C, Ab_D-SC₁, or Ab_E is employed, each different type preferably has a different K.

The amount of L-Ab_C employed can be any amount sufficient to extend the commencement of the hook effect and up to that which gives unsatisfactory non-specific adsorption. More particularly, this amount can range from about 0.01 to about 1 µg per ml of reagent.

The amount of Ab_D employed can be any amount sufficient to extend the commencement of the hook effect and up to the amount required to saturate SC₁. More particularly, this amount can range from 0.1 to about 10 µg per test.

The amount of Ab_E employed can be any amount sufficient to extend the commencement of the hook effect.

In order to obtain the advantages of the present invention in a one- or two-step sandwich immunoassay, one need only employ either L-Ab_C or Ab_D-SC₁. Since it is easier to adjust the labeled antibody concentration and since it is also easier to adjust the ratio of L-Ab_A to L-Ab_C, it is preferred to only employ L-Ab_C as the additional entity.

Ab_A, Ab_B, Ab_C, Ab_D, and Ab_E are each independently selected from a group consisting of monoclonal antibodies, polyclonal antibodies, and mixtures thereof. For example, in one- and two-step sandwich immunoassays, (a) Ab_A, Ab_B, Ab_C, and Ab_D can be monoclonal antibodies or can be polyclonal antibodies; or (b) Ab_A and Ab_C can be both monoclonal antibodies and Ab_B and Ab_D can be both polyclonal antibodies; or (c) Ab_A and Ab_C can be both polyclonal antibodies and Ab_B and Ab_D are

both monoclonal antibodies. In one- and two-step sandwich immunoassays, Ab_c and Ab_d are preferably monoclonal antibodies.

The following examples are provided for purposes of further illustration only and are not intended to be limitations on the disclosed invention.

Example 1

IgE Enzyme Immuno Assay (Prior Art)

Materials

Polystyrene bead coated with a monoclonal antibody directed against a specific IgE site (Ab_b -SC).

Horse radish peroxidase-labeled antibody directed against different IgE sites (L - Ab_a) and having a K of about 2×10^{10} .

o-Phenylenediamine

Hydrogen peroxide

Stopping solution

IgE Standards 0, 10, 25, 75, 200, and 400 IU/ml

Substrate buffer

Protocol

1. Allow all components and samples to equilibrate to room temperature and mix well prior to use.
2. Add 20 μ l standard (including zero dose or sample to each test tube.

3. Add 300 μ l diluted conjugate to each tube.
4. Mix by gently shaking tube rack. Tap rack to displace air bubbles.
5. Remove bead basket from container with forceps. Hold over container until drained and place on inverted cap. Place one bead in each tube.
6. Gently shake tube rack to mix. Tap rack to remove air bubbles if necessary.
7. Incubate the tubes in a $37 \pm 1^\circ$ C. water bath for 30 minutes.
8. Prepare the substrate solution during the last five minutes of the incubation. Use plastic forceps to handle tablets.
9. Wash beads by filling each tube with approximately 3 ml distilled water and aspirate. Repeat two more times.
10. At timed intervals, add 300 μ l of enzyme substrate solution to each tube.
11. Shake test tube rack to ensure mixing and cover tubes to exclude light.
12. Incubate 30 ± 1 minute at 20° C. to 25° C.
13. At timed intervals, add 1 ml Stopping Solution to each tube. Add in the same sequence and with the same timing as used for substrate addition. Mix by gentle shaking.

14. Zero the spectrophotometer against distilled water and measure the absorbance of samples and standards at 492 nm. Calculate corrected absorbances for each standard and sample by subtracting the absorbance of the reagent blank (zero dose).

Calculations

1. On semi-logarithmic graph paper, label the linear axis with absorbance (0 to 2.00A) and label the logarithmic axis with the concentration of the standards (10 to 400 IU/ml).
2. Prepare the standard curve by plotting the average corrected absorbances of the standards, and connecting the points with straight lines.
3. Use the corrected absorbances of the samples to interpolate sample concentrations from the standard curve.

The results of this experiment are set forth in Table I.

Example 2

IgE Enzyme Immunoassay (Present Invention)

The materials, protocol, and calculations set forth in Example 1 were employed with one modification. The sole modification consisted of the use of one additional type of horse radish peroxide-labeled antibody directed against different IgE sites (L-Ab_C) and having a K of about 3×10^9 . The results of this experiment are also set forth in Table I.

TABLE I

<u>IgE, IU/ml</u>	<u>Absorbance</u>	
	<u>Prior Art</u> <u>(Example 1)</u>	<u>Present Invention</u> <u>(Example 2)</u>
0	0.089	0.126
10	0.161	0.186
25	0.282	0.323
75	0.614	0.634
200	1.069	1.099
400	1.396	1.592
1,000	1.916	2.226
4,000	1.994	2.808
8,000	1.551	2.718
16,000	1.387	3.018
40,000	1.013	2.826

The IgE assays of Examples 1 and 2 have a dynamic range of from 0 to 400 IU IgE per ml. Accordingly, the data set forth in Table I indicate that at IgE concentrations greater than 8000 IU/ml, one would obtain a false negative result with the prior art procedure because of the presence of the hook effect. In contrast, with the procedure and kit of the present invention, the hook effect is avoided for IgE concentrations of at least 40,000 IU/ml. As the data of Table I also show, this avoidance is accomplished without any significant elevation of signal in the dynamic range.

Example 3HCG Enzyme Immuno Assay (Prior Art)Materials

Polystyrene bead coated with a monoclonal antibody directed against a specific HCG site (Ab_b-SC).



Horse radish peroxidase-labeled antibody directed against different HCG sites (L-Ab_a) and having a K of about 8×10^{10} .

o-Phenylenediamine

Hydrogen peroxide

Stopping solution

HCG Standards 0, 10, 25, 75, 200, and 400 IU/ml

Substrate buffer

Protocol

1. Allow all components and samples to equilibrate to room temperature and mix well prior to use.
2. Add 200 μ l standard (including zero dose or sample to each test tube.
3. Add 200 μ l diluted conjugate to each tube.
4. Mix by gently shaking tube rack. Tap rack to displace air bubbles.
5. Remove bead basket from container with forceps. Hold over container until drained and place on inverted cap. Place one bead in each tube.
6. Gently shake tube rack to mix. Tap rack to remove air bubbles if necessary.
7. Incubate the tubes on a clinical rotor at 190 ± 10 RPM for 45 minutes at 20° C. to 25° C.

8. Prepare the substrate solution during the last five minutes of the incubation. Use plastic forceps to handle tablets.
9. Wash beads by filling each tube with approximately 3 ml distilled water and aspirate. Repeat two more times.
10. At timed intervals, add 300 μ l of enzyme substrate solution to each tube.
11. Shake test tube rack to ensure mixing and cover tubes to exclude light.
12. Incubate 30 ± 1 minute at 20° C. to 25° C.
13. At timed intervals, add 1 ml Stopping Solution to each tube. Add in the same sequence and with the same timing as used for substrate addition. Mix by gentle shaking.
14. Zero the spectrophotometer against distilled water and measure the absorbance of samples and standards at 492 nm. Calculate corrected absorbances for each standard and sample by subtracting the absorbance of the reagent blank (zero dose).

Calculations

1. On semi-logarithmic graph paper, label the linear axis with absorbance (0 to 2.00A) and label the logarithmic axis with the concentration of the standards (1 to 100 mIU/ml).



2. Prepare the standard curve by plotting the average corrected absorbances of the standards, and connecting the points with straight lines.
3. Use the corrected absorbances of the samples to interpolate sample concentrations from the standard curve.

The results of this experiment are set forth in Table II.

Example 4

HCG Enzyme Immuno Assay (Present Invention)


The materials, protocol, and calculations set forth in Example 3 were employed with one modification. The sole modification consisted of the use of one additional type of horse radish peroxide-labeled antibody directed against different HCG sites (L-Ab_C) and having a K of about 2×10^9 . The results of this experiment are also set forth in Table II.

TABLE II

HCG, mIU/ml	Absorbance	
	Prior Art (Example 3)	Present Invention (Example 4)
0	0.090	0.092
1	0.101	0.108
2.5	0.130	0.132
5	0.173	0.188
10	0.290	0.284
25	0.546	0.557
50	0.952	0.932
100	1.620	1.592
1000	3.82	3.855
10000	1.64	5.514
64880	0.66	4.37
128000	0.42	3.38

The HCG assays of Examples 3 and 4 have a dynamic range from 0 to 100 mIU HCG per ml. Accordingly, the data set forth in Table II indicate that at HCG concentrations greater than 10.000 mIU/ml, one would obtain a false negative result with the prior art procedure because of the presence of the hook effect. In contrast, with the procedure and kit of the present invention, the hook effect is avoided for HCG concentrations of at least 128,000 mIU/ml. As the data of Table II also show, this avoidance is accomplished without any significant elevation of signal in the dynamic range.

Based on this disclosure, many other modifications and ramifications will naturally suggest themselves to those skilled in the art. These are intended to be comprehended as within the scope of this invention.



Bibliography

1. U.S. 4,376,110
2. U.S. 4,244,940
3. Miles et al., Analytical Biochemistry, 61:209-224 (1974)
4. Uotila et al., Journal of Immunological Methods, 42:11-15 (1981)
5. Ng et al., Clin. Chem., 29(6):1109-113 (1953)
6. Nomura et al., Journal of Immunological Methods, 56:13-17 (1983)
7. Miles et al., Symposium on Radioimmunoassay and Related Procedures in Clinical Medicine and Research, International Atomic Energy Agency, Vienna, Austria (1973) pp. 149-164.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. An improved immunoassay for an antigenic substance (Ag) in a fluid, said immunoassay being of the type comprising contacting said fluid with at least one first entity selected from a group consisting of (a) an antibody (Ab) to said Ag, (b) soluble, labeled antibody (L-Ab_a) to said Ag, and (c) antibody (Ab_b) to said Ag bound to a solid carrier (SC), characterized in that said fluid is contacted with at least one additional entity selected from a group consisting of at least one different type of soluble labeled antibody (L-Ab_c), at least one different type of antibody (Ab_d) to said Ag bound to a solid carrier (SC₁), and at least one different type of antibody (Ab_e) to said Ag,

wherein:

- (i) each of said additional entities has an average affinity constant (K) for said Ag lower than the K of its corresponding first entity for said Ag;
 - (ii) said additional entity is present in an amount sufficient to avoid a hook effect; and
 - (iii) said SC₁ is selected from a group consisting of said SC, at least one different solid carrier (SC₂), and mixtures thereof (SC and SC₂).
2. The improved immunoassay of claim 1 wherein said immunoassay is a one-step sandwich immunoassay of the type comprising:
 - (a) contacting said fluid with (i) said soluble, labeled antibody (L-Ab_a) to said Ag and (ii)



said antibody (Ab_b) to said Ag bound to said solid carrier (SC) to form an insoluble complex ($L-Ab_a-Ag-Ab_b-SC$);

(b) separating said $L-Ab_a-Ag-Ab_b-SC$ from said fluid and unreacted $L-Ab_a$; and

(c) measuring either the amount of $L-Ab_a$ associated with said $L-Ab_a-Ag-Ab_b-SC$ or the amount of unreacted $L-Ab_a$;

and is characterized in that said fluid is contacted with at least one additional entity selected from said group consisting of at least one different type of soluble, labeled antibody ($L-Ab_c$) to said Ag and at least one different type of said antibody (Ab_d) to said Ag bound to said solid carrier (SC_1);

wherein:

(i) each type of said $L-Ab_c$ and Ab_d-SC_1 has a lower average affinity constant (K) for said Ag than each respective K of said $L-Ab_a$ and said Ab_b-SC .

3. The improved immunoassay of claim 2 wherein said additional entity is said $L-Ab_c$.
4. The improved immunoassay of claim 2 wherein Ab_a , Ab_b , Ab_c , and Ab_d are each independently selected from a group consisting of monoclonal antibodies, polyclonal antibodies, and mixtures thereof.
5. The improved immunoassay of claim 2 wherein Ab_a and Ab_c are monoclonal antibodies.
6. The improved immunoassay of claim 4 wherein said additional entity is said $L-Ab_c$.

7. The improved immunoassay of claim 1 wherein said immunoassay is a two-step sandwich immunoassay of the type comprising:
- (a) contacting said fluid with said antibody (Ab_D) bound to a solid carrier (SC) to form an insoluble complex ($SC-Ab_D-Ag$);
 - (b) contacting said $SC-Ab_D-Ag$ with a soluble labeled antibody complex ($SC-Ab_D-Ag-Ab_A-L$);
 - (c) separating said $SC-Ab_D-Ag-Ab_A-L$ from unreacted $L-Ab_A$; and
 - (d) measuring either the amount of $L-Ab_A$ associated with said $SC-Ab_D-Ag-Ab_A-L$ or the amount of unreacted $L-Ab_A$;
- and is characterized in that in step (b) said $SC-Ab_D-Ag$ is contacted with at least one different type of soluble, labeled antibody ($L-Ab_C$);
- wherein:
- (i) each type of said $L-Ab_C$ has a lower average affinity constant (K) for said Ag than the K of said $L-Ab_A$.
8. The improved immunoassay of claim 6 wherein Ab_A , Ab_D , and Ab_C are each independently selected from a group consisting of monoclonal antibodies, polyclonal antibodies, and mixtures thereof.
9. The improved immunoassay of claim 7 wherein Ab_A and Ab_C are monoclonal antibodies.
10. The improved immunoassay of claim 1 wherein said immunoassay is a direct nephelometric immunoassay of the type comprising:
- (a) contacting said fluid with said antibody (Ab) to said Ag in order to form a complex $Ab-Ag$; and
 - (b) measuring the amount of formed $Ab-Ag$;

and is characterized in that said fluid is contacted with at least one additional type of antibody Ab_e , each type of said Ab_e having an average affinity constant (K) for said Ag lower than the K of said Ab for said Ag.

11. The improved direct nephelometric immunoassay of claim 10 wherein each additional type of Ab_e is a monoclonal antibody.
12. The improved direct nephelometric immunoassay of claim 16 wherein each additional type of Ab_e is a polyclonal antibody.
13. An improved reagent of the type comprising at least one first entity selected from a group consisting of (a) an antibody (Ab) to an antigenic substance (Ag); (b) a soluble, labeled antibody ($L-Ab_a$) to said Ag; and (c) an antibody (Ab_b) bound to a solid carrier (SC), characterized in that said reagent further comprises at least one additional entity selected from a group consisting of at least one different type of soluble labeled antibody ($L-Ab_c$), at least one different type of antibody (Ab_d) to said Ag bound to a solid carrier (SC_1), and at least one different type of antibody (Ab_e) to said Ag, wherein:
 - (i) each of said additional entities has a lower average affinity constant (K) for said Ag than the K of its corresponding first entity;
 - (ii) said additional entity is present in an amount sufficient to avoid a hook effect when said reagent is employed in an immunoassay; and

- (iii) said SC_1 is selected from a group consisting of said SC, at least one different solid carrier (SC_2), and mixtures thereof (SC and SC_2).

14. The improved reagent of claim 13 being of the type comprising:
- (a) said $L-Ab_a$ to said Ag; and
 - (b) said Ab_b bound to said SC;
- and being characterized in that said reagent further comprises at least one additional entity selected from a group consisting of at least one different type of soluble labeled antibody ($L-Ab_c$) to said Ag and at least one different type of antibody (Ab_d) to said Ag bound to a solid carrier (SC_1); wherein:
- (i) each type of said $L-Ab_c$ and said Ab_d-SC_1 has a lower K for said Ag than each respective K of said $L-Ab_a$ and said Ab_b-SC .
15. The improved reagent of claim 14 comprising:
- (a) a first unit comprising said $L-Ab_a$ to said Ag (and said $L-Ab_c$ to said Ag) $_x$; and
 - (b) a second unit comprising said Ab_b-SC (and said Ab_d-SC_1) $_y$;
- where x and y are 0 or 1, provided that when x and y are equal, x and y are 1.
16. The improved reagent of claim 15 wherein x is 1 and y is 0.
17. The improved reagent of claim 7 wherein said Ab_a , Ab_b , Ab_c , and Ab_d are each independently selected from a group consisting of monoclonal antibodies, polyclonal antibodies, and mixtures thereof.

18. The improved reagent of claim 17 wherein said Ab_a and Ab_c are monoclonal antibodies.
19. The improved reagent of claim 13 being of the type comprising said Ab , and being characterized in that said reagent further comprises at least one different type of Ab_e , each different type of said Ab_e having a K for said Ag lower than the K of said Ab for said Ag.
20. The improved reagent of claim 19 wherein each of said Ab_e is independently selected from a group consisting of monoclonal antibodies, polyclonal antibodies, and mixtures thereof.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 84/01888

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁴ : G 01 N 33/53; G 01 N 33/543; G 01 N 33/577		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System ¹	Classification Symbols	
IPC ⁴	G 01 N	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	Molecular Immunology, vol. 17, published in 1980, Pergamon Press Ltd. (GB) J. Steensgaard et al.: "The development of difference turbidimetric analysis for monoclonal antibodies to human IgG", see pages 1315-1318 --	
A	Clinical Chemistry, vol. 27, no. 11, published in November 1981 (Easton, Pennsylvania, US) E.D. Sevier et al.: "Monoclonal antibodies in clinical immunology", see pages 1797-1806 --	
A	EP, A, 0045103 (AKZO N.V.) 3 February 1982 --	
A	EP, A, 0048357 (LA JOLLA CANCER RESEARCH FOUNDATION) 31 March 1982 --	
A	EP, A, 0028133 (DYNASCIENCES CORPORATION) 6 May 1981 --	./.
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
27th March 1985	17 AVR. 1985	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	G.L.M. Kruyenberga	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	EP, A, 0062892 (AMERICAN HOECHST CORP.) 20 October 1982 -----	

INTERNATIONAL APPLICATION NO.

PCT/US 8401888 (SA 8420)

This Annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 12/04/85

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0045103	03/02/82	NL-A- 8004308	01/03/82
		AU-A- 7307891	04/02/82
		JP-A- 57086051	28/05/82
EP-A- 0048357	31/03/82	JP-A- 57079455	18/05/82
		SE-A- 8006424	13/03/82
		CA-A- 1171780	31/07/84
EP-A- 0028133	06/05/81	JP-A- 56067757	08/06/81
		AU-A- 6295180	30/04/81
		CA-A- 1146856	24/05/83
EP-A- 0062892	20/10/82	JP-A- 57167653	15/10/82
		US-A- 4468856	04/09/84